

Synthesis, spectral studies and biological applications of benzoic acid and its derivatives

Muhammad Aqeel Ashraf; Mohd. Jamil Maah; Ismail Yusoff

Department of chemistry, University of Malaya, 50603 Kuala Lumpur, Malaysia
chemaqeel@gmail.com

Abstract—Resistance to the presently accessible antibiotics has motivated the search for new agents to inhibit bacterial activity. Among such agents, metal complexes of biologically active ligands are attractive as both ligands and metal ions can interact with different steps of pathogenic life cycles. Much work has been done by bioinorganic as well as medicinal chemists to establish the relationship between the metal ions and their complexes as antitumor and antibacterial agents. In this study Cr (III), Co (II), Ni (II), Cu (II) and Zn (II) salts of 2-(4-nitrophenylaminocarbonyl) benzoic acid have been prepared and were characterized by physical, analytical and spectroscopic studies. On the basis of electronic spectra and electronic susceptibility measurements in conjunction with infrared spectra, six coordinated octahedral structure have been proposed to all the complexes. The benzoic acid and these complexes have been tested for their *in-vitro* antimicrobial activity against three bacterial strains, *Mycobacterium smegmatis* (Gram +ve), *Escherichia coli* (Gram -ve), *Pseudomonas aeruginosa* (Gram -ve) and three fungal strains, *Nigrospora oryzae*, *Aspergillus niger* and *Candida albicans*. The antimicrobial activities of the metal complexes were found to be greater than those of 2-(4-nitrophenylaminocarbonyl) benzoic acid alone.

Keywords—4-nitroaniline; Phthalic anhydride; Metal ions; Antimicrobial activity.

I. INTRODUCTION

Coordination compounds have been a challenge to inorganic chemist since they were identified in the 19th century. They defy the usual rules of valence at that time and hence called complexes. They play vital role in our lives. Transitions metal complexes with soft or hard donor groups have been used extensively in coordination and organometallic chemistry [1]. Reaction of coordinated ligands model systems of biological interest, analytical chemistry, agriculture and optical display devices [2-4]. Resistance to the presently accessible antibiotics has motivated the search for new agents to inhibit bacterial activity. Among such agents, metal complexes of biologically active ligands are attractive as both ligands and metal ions can interact with different steps of pathogenic life cycles[5-7]. Much work has been done by bioinorganic as well as medicinal chemists to establish the relationship between the metal ions and their complexes as antitumor and antibacterial agents [8-12]. It is however remarkable that some biologically active compounds may become more carcinostatic and bacteriostatic upon chelation [13]. There is

currently a resurgence of interest in the biochemistry as well as the coordination chemistry of bivalent Cr, Co, Ni, Cu and Zn due to their biological importance.

In view of this and our interest in biologically active coordination compound, the present work communicating the synthesis, characterization and biological activity of M^{2+} ($M=Cr, Co, Ni, Cu, Zn$) with benzoic acid as ligand. Aniline derivatives have been actively investigated because of their fascinating biological and diverse ligational behaviour towards metal ions and novel structural features in their complexes [14-25]. We have made studied the coordination and biological chemistry of 2-(4-nitrophenylaminocarbonyl)benzoic acid.

Polluted water consists of Industrial discharged effluents, sewage water, and the rain water. The use of this type of water is a common practice in agriculture. Estimation indicates that more than 50 countries of the world with an area of twenty million hectares area are treated with polluted or partially treated polluted water [1]. In poor countries of the world more than 80% polluted water have been used for irrigation with only seventy to eighty percent food and living security in industrial urban and semi urban areas [2].

II. MATERIAL AND METHODS

All chemicals used were of analytical grade and were obtained from Merck. I.R spectra were recorded on a Spectrum 100 FTIR Spectrophotometer Perkin Elmer USA. U.V-Visible spectra were obtained in DMSO by Agilent 8453-Spectrophotometer using ChemStation software. Melting points were determined by a melting point meter M5000 from KRUSS Optronic.

A. Synthesis of Ligand

2-(4-nitrophenylaminocarbonyl)benzoic acid was made as described previously[19, 35].




Figure 1. Synthesis of 2-(4-nitrophenylaminocarbonyl)benzoic acid

B. Synthesis of Metal Complexes

• Synthesis of Cu (II) – Complex

2-(4-nitrophenylaminocarbonyl) benzoic acid (0.01 mol, 0.286 g) was suspended in acetone (100 mL) in a 250 mL two necked round bottom flask and then treated with triethylamine (0.45 mL). The mixture was refluxed for 2-3 h. Then $\text{CuCl}_2 \cdot 2\text{H}_2\text{O}$ (0.005 mol) was added to the reaction flask with constant stirring and the reaction mixture was again refluxed for 8–10 h. A dark green precipitate was formed. The solvent was evaporated and the remaining mass was recrystallized from CHCl_3 / n-hexane mixture (1:1) and was stored in a moisture-free environment.

The Ni(II), Cr(II), Co(II) and Zn(II) complexes were made similarly from the hydrated chlorides.




Figure 2. Proposed structures of metal complexes

III. BIOLOGICAL STUDIES

A. Antibacterial studies

The ligand as well as its metal complexes were screened against three bacterial species ; *Mycobacterium smegmatis*, *Escherichia coli*, *Pseudomonas aeruginosa* by using the paper disc diffusion method [17], nutrient agar medium (Peptone, Beef extract, NaCl and Agar-Agar) and 5mm diameter paper discs (Whatman No.1). The compounds *i.e.* ligand precursor and its complexes, were dissolved (30 μg) in DMF (0.01 mL). The filter paper disc were soaked in these solutions, dried, and then placed in Petri plates previously seeded with the test organisms. The plates were incubated for 24 h at 37°C and the inhibition zone in the region of each disc was measured.

B. Antifungal Activities

The antifungal activity of the ligand and its metal complexes were evaluated against *Nigrospora oryzae*, *Aspergillus niger* and *Candida albicans* by the agar plate technique using the reported method [36].

IV. RESULTS AND DISCUSSION

A. Physical analysis

2-(4-Nitrophenylaminocarbonyl)benzoic acid reacted with solutions of metal ions to give colored amorphous complexes of general formulae ML_2 and $\text{ML}_2(\text{H}_2\text{O})_2$ as shown by elemental analysis (Table-3).

They were insoluble in common organic solvents and soluble in DMF and DMSO[23].

B. Electronic spectral analysis

Data obtained from the electronic spectroscopy are tabulated in Table-1. For the chromium complex two peaks at 15898 and 23770 cm^{-1} were assigned to ${}^4A_{2g} \rightarrow {}^4T_{2g}$ and ${}^4A_{2g} \rightarrow {}^4T_{1g}(f)$ d-d transitions, respectively. The appearance of these two bands confirms octahedral (O_h) geometry for this complex. The electronic spectrum of the Cu(II)-complex showed bands at 31055 cm^{-1} and 19820 cm^{-1} and are assigned, respectively, to charge transfer and $E_g \leftarrow {}^2T_{2g}$ transitions. These two band are in agreement with those usually observed for square planar Cu(II)-complexes [26]. The electronic spectrum of the Co(II) complex showed bands at 7373 cm^{-1} , 17213 cm^{-1} and 20613 cm^{-1} assigned to ${}^4T_{1g}(F) \rightarrow {}^4T_{2g}(F)$, ${}^4T_{1g}(F) \rightarrow {}^4A_{2g}(F)$ and ${}^4T_{1g}(F) \rightarrow {}^4T_{2g}(P)$ transitions, respectively, corresponding to octahedral geometry. The Ni(II)-complex also exhibited three spin-allowed bands at 26312 cm^{-1} , 15433 cm^{-1} , and 9789 cm^{-1} , assigned, respectively, to ${}^3A_{2g}(F) \rightarrow {}^3T_{2g}(F)$, ${}^3A_{2g}(F) \rightarrow {}^3T_{1g}(F)$, and ${}^3A_{2g}(F) \rightarrow {}^3T_{2g}(P)$ transitions, consistent with well defined octahedral geometry. The diamagnetic Zn(II)-complex did not show any d-d transition.

Table 1: Physical Properties of Ligand and its Metal Complexes

S.No	Compound	Color	Physical State	Melting Point (°C)	μ_B (B.M)	Conductance $\mu\text{S}/\text{cm}$	Yield %	λ_{max} (cm^{-1})
1.	Ligand	Yellowish brown	Crystalline	230	-	-	75	-
2.	Cr(III)-complex	Dark Green	Amorphous	-280	3.81	121	68	15898, 23770
3.	Ni(II)-complex	Brownish Yellow	Amorphous	-290	3.12	19	70	6312, 15433, 9789
4.	Co(II)-complex	Light Pink	Amorphous	-291	4.76	33	56	7373, 17213, 20613
5.	Cu(II)-complex	Green	Amorphous	-285	1.79	12	62	31055, 19820
6.	Zn(II)-complex	White	Amorphous	-267	Diamagnetic	14	43	27123

Ligand = 2-(4-nitrophenylaminocarbonyl)benzoic acid

C. Magnetic moment

The magnetic moment value for the Cr(III)-complex was found to be 3.81 BM (table-1) which suggests an octahedral geometry for this complex [27]. The magnetic moment for value Cu(II)-complex (1.79 BM) also favors its square planar geometry. The magnetic moment value (3.12 BM) showed two unpaired electron per Ni (II) ion and is indicative of octahedral environment for Ni(II). Similarly the magnetic moment of the Co (II)-complex (4.76 BM) is consistent with octahedral geometry [27]. The Zn(II) complex was found to be diamagnetic

Table 2: IR stretching frequencies (cm⁻¹) of the various functional groups of ligand and its metal complexes

S.No	Compound	C-O cm ⁻¹	NH cm ⁻¹	(COO) asym cm ⁻¹	(COO) Sym cm ⁻¹	Δν cm ⁻¹	(OH) cm ⁻¹	C-H cm ⁻¹	N-O asym. cm ⁻¹	H ₂ O cm ⁻¹	C-O-C cm ⁻¹	C-N cm ⁻¹
1.	A	-	3479	-	-	-	-	794	1504	-	-	-
2.	B	1762	-	-	-	-	-	798	-	-	1280	-
3.	Ligand	1702	3379	1595	1377	218	2853	798	1518	-	-	1306
4.	Cr(III)- complex	1730	3373	1553	1415	138	-	794	1525	3667	-	1295
5.	Ni(II)-complex	1729	3378	1566	1381	185	-	709	1519	3643	-	1300
6.	Co(II)-complex	1728	3376	1550	1378	172	-	783	1521	3455	-	1308
7.	Cu(II)-complex	1729	3378	1552	1384	168	-	708	1514	-	-	1290
8.	Zn(II)-complex	1732	3370	1559	1376	183	-	722	1513	-	-	1298

A = 4-Nitroaniline B = Phthalic Anhydride - = absent

Ligand = 2-(4-nitrophenylaminocarbonyl)benzoic acid

D. IR spectral analysis

The IR stretching frequencies and assignments are given in table 2.

A carboxylate ligand can bind to the metal in monodentate or bidentate fashion, resulting in changes in the relative positions of the asymmetric and symmetric stretching vibrations [28]. The IR spectra of the complexes give a separation value ($\Delta\nu$) less than 200 cm⁻¹, which confirms that the coordination is bidentate [29-30]. A strong band at 3479-3437 cm⁻¹, characteristic of the NH group and present in the spectrum of the ligand precursor persists in the spectra of all metal complexes. This shows that NH group does not participate via intra or intermolecular interactions [31]. Peaks above 3400 cm⁻¹ in the Co(II), Ni(II) and Cr(II) complexes, indicated the presence of coordinated water [17]. These were not present in the Cu(II) and Zn(II) complexes, suggesting that the coordination number in these compounds is only four. All the above discussion is consistent with the structures in Figure-1.

Table 3. Elemental analysis data of ligand and its metal complexes

S.No	Compound	C	H	N	M.
		% Obs. (Cal.)	% Obs. (Cal.)	% Obs. (Cal.)	% Obs. (Cal.)
1	Ligand	58.43 (58.95)	3.07 (3.16)	9.63 (9.82)	-
2	Cr(III)-complex	50.87 (51.09)	2.33 (2.73)	8.23 (8.51)	7.77 (7.90)
3	Ni(II)-complex	50.34 (50.57)	2.57 (2.71)	8.19 (8.42)	8.45 (8.83)
4	Co(II)-complex	50.43 (50.55)	2.67 (2.70)	8.61 (8.41)	8.47 (8.86)
5	Cu(II)-complex	52.87 (53.06)	2.54 (2.84)	8.43 (8.84)	10.56 (11.14)
6	Zn(II)-complex	52.30 (52.90)	2.67 (2.83)	8.57 (8.81)	11.10 (11.47)

Ligand = 2-(4-nitrophenylaminocarbonyl)benzoic acid

E. Biological Studies

a) Antibacterial studies

Metal complexes were screened against three bacterial species like *Mycobacterium smegmatis*, *Escherichia coli*, *Pseudomonas aeruginosa* by the paper disc diffusion method [17]. Studies of *in vitro* antibacterial activity of metal complexes exhibit appreciable antibacterial activity (table-4) that is in accordance with our previous studies [32-33]. This enhanced antibacterial activity may be due to an increase in

cell permeability of the lipophilic metal conjugates, which allows intracellular drug accumulation. It is also likely that the intracellular reduction of these metal compounds may lead to higher cytoplasmic concentration of metal species, which prove lethal for bacteria.

Table 4 : Antibacterial activity of ligand and its metal complexes

S.No	Compound	Bacterial Strain Inhibition Zone (mm)		
		A	B	C
1.	Sulphadimidine	+++	+++	++
2.	Ligand	+	-	+
3.	Cr(III)-complex	++	+	+
4.	Ni(II)-complex	+	++	-
5.	Co(II)-complex	++	++	++
6.	Cu(II)-complex	++	++	+++
7.	Zn(II)-complex	+++	+	++

Ligand = 2-(4-nitrophenylaminocarbonyl)benzoic acid

A = *E. coli* B = *Pseudomonas aeruginosa* C = *Mycobacterium smegmatis*
Inhibition Zone - + 0-5 mm, ++ 6-0 mm, +++ 11-15mm, - inactive.

b) Antifungal studies

The antifungal activity of all the synthesized metal complexes was determined against three fungal strains namely; *Nigrospora oryzae*, *Aspergillus niger* and *Candida albicans* by the agar plate technique and the results are recorded in table-5. It was observed that metal complexes are more active than the corresponding ligand and toxicity increases with increasing concentration [34].

Table 5: Antifungal activity of ligand and its metal complexes

S.No.	Compound	A.			B.			C.		
		100µg/ml.	200µg/ml.	1000µg/ml.	100µg/ml.	200µg/ml.	1000µg/ml.	100µg/ml.	200µg/ml.	1000µg/ml.
1.	Ketoconazole	10.5	15.8	27.9	15.5	23.8	31.9	12.3	17.7	33.8
2.	Ligand	7.2	12.1	21.4	-	2.4	5.6	10.3	15.3	22.2
3.	Cr(III)-complex	10.2	14.3	16.0	17.2	23.0	26.5	19.5	33.6	45.2
4.	Ni(II)-complex	8.0	13.5	25.8	10.2	14.5	19.4	22.4	38.3	56.2
5.	Co(II)-complex	15.2	35.5	54.4	16.2	30.3	44.9	13.3	24.5	34.4
6.	Cu(II)-complex	22.6	47.5	63.4	17.3	27.7	34.2	21.5	31.2	49.4
7.	Zn(II)-complex	21.2	35.5	53.4	15.2	28.3	30.5	13.3	29.5	39.4

Ligand = 2-(4-nitrophenylaminocarbonyl)benzoic acid

a. *Nigrospora oryzae* b. *Aspergillus niger* c. *Candida albicans* - inactive

V. ACKNOWLEDGMENT

The work reported in this paper is a basic research work carried out in Analytical Laboratory, Department of Chemistry, University of Malaya. I take this opportunity to express my gratitude to Ministry of Higher Education, Malaysia to provide me scholarship to complete my higher studies in Malaysia. Thanks also go to Institut Pengurusan Dan Pemantauan Penyelidikan, (IPPP) University of Malaya for providing me enough funding to complete this valuable research.

VI. REFERENCES

- [1] Sarkar, A. R. & Mandal, S. 2000. Synthesis and Reactivity in Inorganic and Metael Organic Chemistry, Vol.30, p.1477.
- [2] Mishra, A. P. & Khare, M. 2000. Journal of the Indian Chemical Society, 77 (8):367-377.
- [3] Siddiqui, R. S. & Nishat, N. 2003. Synthesis and Reactivity in Inorganic and Metael Organic Chemistry, 33 (4):565-589.
- [4] Maurya, M. R. & Bharatia, N. 1993. Transition Metals Chemistry, 23, 5-15.
- [5] Scozzafava, A. & Supuran, C. T. 2000. Journal of medicinal Chemistry, 43: 3677-3687.
- [6] Brown, D. H. Smith, W.E. Teape, J. W & Lewis, A. J. 1980. Journal of medicinal Chemistry, 23: 729-734.
- [7] Scozzafava, A. Menabuoni, L. Mincione, F. Mincione G. & Supuran, C. T. 2001. Bioorganic Medicinal & Chemistry Letters, 11: 575-582.
- [8] William. R. J. P. 1970. Quarterly Review, 24: 331-365.
- [9] Rosenberg, B. & Camp, L. V. 1970. Cancer Research, 30: 1799-1802.
- [10] Clare, M. J. & Heeschele, J.D. 1973. Bioinorganic Chemistry, 2: 187-195.
- [11] Narayanan, V. L. Nasr, M. & Paull, K. D.1990. Tin-based Antitumour drugs, p 200, M. Gielen (Ed), NATOASI Series: Springer: Berlin.
- [12] Crowe, A. J. 1988. Metal based antitumour drugs, p 103, M. Gielen (Ed), Freund: London.
- [13] Srivastava, R. S. Indian Journal of Chemistry. 29: 1024-1026.
- [14] Marteil, A. E. 1968. Pure and Applied Chemistry, 12: 129-134.
- [15] Khuhawar, M. Y. Journal of Chemical Society of Pakistan, 6: 225-231.
- [16] Lieno, K. & Martell, A. E. 1957. Journal of Physical Chemistry, 41: 257-263.
- [17] Imran, M. Iqbal, J. & Latif, S. 2008. Journal of Chemical Society of Pakistan, 30: 594-598.
- [18] Iqbal, J. Tirmizi, S. A. Wattoo, F. H. Imran, M. Wattoo, H. S. Sharfudin, S. & Latif, S. 2006. Turkish journal of Biology, 30: 1-4.
- [19] Shahid, K. Ali, S. Shahzadi, S. & Akhtar, Z. 2003. Turkish Journal of Chemistry, 27: 209-215.
- [20] Shahid, K. Shahzadi, S. Ali, S. Bhatti, M. H. & Khan, K. M. 2005. Journal of Iranian Chemical Society, 2: 140-148.
- [21] Park, J. O. & Jang, S. H. 1992. Journal of Polymer Science Part A Polymer Chemistry, 30: 723-729.
- [22] Shahzadi, S. Shahid, K. Ali, S. & Bakhtiar, M. 2008. Turkish Journal of Chemistry, 32:333-340.
- [23] Hassan, M. Scozzafava, A. Chohan, Z. H. & Supuran, C. T. 2003. Journal of Enzyme Inhibition and Medicinal Chemistry, 18: 495-503
- [24] Parashar, R. K. Sharma, R. C. & Mohan, G. 1990. Biological Trace Element Research, 23: 150-158.
- [25] Phatak, P. Jolly, V. S. & Sharma, K. P. 2000. Oriental Journal of Chemistry, 16: 493-494.
- [26] Lever, A. B. 1968. Organic electronic Spectroscopy, p 307, 2nd ed., Elsevier: Amsterdam.
- [27] Cotton, F. A. Wilkinson, G. Murillo, C. A. & Bochmann, M. 1999. Advanced inorganic chemistry, p 821, 6th Ed, John Wiley and Sons: New York.
- [28] Xie, L. Q. Yang, Z. Q. Zhang, Z. X. & Zhang, D. K. 1992. Applied Organometallic Chemistry, 6: 193-199.
- [29] Xie, Q. Yang, Z. and Jiang, L. 1996. Main Group Metal Chemistry, 19: 509-520.
- [30] Imran, M. Iqbal, J. Iqbal, S. & Latif, S. 2006. Journal of Chemical Society of Pakistan, 29: 151-155.
- [31] Kemp, W. Organic spectroscopy, 1991. p 79, 3rd Ed, Palgrave; New York.
- [32] Imran, M. Iqbal, J. Mehmood, T & Latif, S. 2006. Journal of Biological Sciences, 6: 946-949.
- [33] Imran, M. Iqbal, J. Iqbal, S. & Ijaz, N. 2007. Turkish Journal of Biology, 31: 67-72.
- [34] Mishra, L. & Singh, V. K. 1997. Indian Journal of Chemistry, 32: 446-452.
- [35] Meyer & Leaders, 1903. Anals of Physics, 327: 52-58.
- [36] Hassan, M. U. Chohan Z. H. & Supuran, C. T. 2002. Main Group Metal Chemistry, 25: 291-296.