

Relationship between serum adiponectin with anthropometrical and lipid profile biochemical indexes in obese adult men

*Mojtaba Eizadi, Davood Khorshidi, Hussein Doali

Islamic Azad University, Saveh Branch

izadimojtaba2006@yahoo.com

Abstract. Background and objective: This study was conducting aimed at comparing adiponectin serum levels in obese male adults with their peer normal weight subjects as well as determining the relationship between anthropometrical and lipid profile indices and serum adiponectin in obese men. **Material and methods:** Body mass index and other anthropometrical variables as well as concentrations of glucose, triglyceride, total cholesterol and low and high density lipoprotein were measured in 45 obese male adults after 12 to 14 hours overnight fasting and their correlation with serum adiponectin levels were determined. Also, fasting serum adiponectin levels were measured in 35 male adults with normal weight. T statistical method was used to compare adiponectin levels of obese group with normal weight group and regression analysis was used in SPSS software version 15 environment to determine the relationship of adiponectin with each anthropometrical and biochemical variables in obese subjects ($P \leq 0.05$). **Results:** adiponectin Serum level in obese subjects were significantly lower than those with normal weight ($P = 0.021$). Adiponectin level had a significant correlation with total cholesterol, low density lipoprotein, glucose, and systolic and diastolic blood pressure and age of subjects ($P \leq 0.05$). **Conclusion:** The significant relationship between adiponectin and some cardiovascular risk factors in this study supports the hypothesis that reduction of systemic levels of this hormone is associated with some cardiovascular risk factors.

Keywords: cardiovascular risk factors, adiponectin, obesity

1. Introduction

However the biological basis for these processes is not fully recognized yet. White adipose tissue as a huge fat reservoir source is also very important in energy homeostasis. So that it reserves body fat as triglyceride (TG) and drops them into the blood stream in the form of free fatty acids (FFA) when needed (1). Moreover, adipose tissue also plays a role as an endocrine organ in the secretion of some biological activators of energy metabolism called adipocytokines such as leptin, resistin, ghrelin and adiponectin. Meanwhile, adiponectin is a protein secreted from adipose tissue which plays an important role in hyperglycemia and inflammatory mechanisms (2).

Adiponectin levels are correlated inversely with anthropometrical indices of obese subjects and insulin resistant. A Russian study showed that the adiponectin which is independent of other blood factors, play an important role in plasma TG levels. Yamamota et al. study revealed that the relationship between adiponectin and TG, LDL and HDL will be maintained even after matching age, sex and BMI, (3). But unlike the above findings, the statistical result of a recent study showed that despite the direct relationship of adiponectin with HDL, there is no relationship between this peptide hormone and other body weight indices, total cholesterol (TC) TG LDL, insulin resistance and anthropometrical indices. Absence of any relationship between serum adiponectin levels and body fat levels has been also observed in some other studies (4). A recent study showed that the significant relationship of adiponectin with lipid profile indices (TG, TC, LDL, and HDL) is independent of BMI. The study of Ram et al. also showed that there is no significant correlation between fasting adiponectin levels with abdominal fat volume (5). Some other studies also suggest that the relationship of adiponectin with biochemical indicators of lipid profile (TG, TC, LDL, and HDL) is

independent of body adipose tissue levels (6). Reviewing the research findings indicates contradiction in the findings regarding the relationship between adiponectin and lipid profile indicators and anthropometrical variables and there is still no general consensus regarding the relationship of this anti-inflammatory hormone with the mentioned indicators. Hence, the present study was also conducted to determine the relationship between the mentioned indicators in obese male adults.

2. Methodology

Forty five obese male adults ($BMI \geq 30$) with age range of 37 to 48 years old participated in this study randomly. Study inclusion criteria include non-athletic, non-smoker individuals with no chronic disease such as diabetes, asthma and kidney disease and no orthopedic abnormalities. Not participating in regular sports activities within at least last six months was also another study inclusion criterion. 31 non-obese male adults with normal weight ($20 \leq BMI \leq 25$) with similar age conditions as the obese group also participated in the study in order to compare the baseline levels of serum adiponectin in the studied obese subjects with that of normal subjects. To run this study, anthropometrical indices in subjects were measured first. So that Weight of the studied subjects was measured in minimum clothing using made in Taiwan digital height meter scale with 100 grams accuracy. Subjects' height was measured while barefoot using height meter scale. Body mass index (BMI) was calculated through dividing body weight (kilograms) by squared height (square meters). Abdominal circumference and hip circumference were measured in the most condensed part using a non-elastic cloth meter. Also, Waist to hip circumference ratio (WHO) was calculated through dividing the abdominal circumference by hip circumference. Subjects were asked to avoid doing any heavy physical activity for 48 hours before blood sampling. The studied subjects were requested to come to the Hematology Laboratory between 8 to 9 am after 12 to 14 hours overnight fasting in order to measure biochemical variables. Then, 8 cc bloods were taken from the subjects' brachial vein to measure fasting adiponectin and lipid profile indicators including (TG, TC, LDL, and HDL). Glucose level of fasting blood was measured through glucose oxidase enzymatic method (Pars Azmoon kit). Independent T-test was used to compare the baseline serum levels of adiponectin in obese group with that of subjects with normal weight. Linear regression test was also used in version 15 SPSS statistical software environment to determine the relationship between serum adiponectin levels with each anthropometrical indices or biochemical variables of lipid profile. In this study, the significance level was considered equivalent to $p < 0.05$ (Saveh, summer 1389).

3. Results

Mean and standard deviation of anthropometrical characteristics, biochemical variables as well as significance levels of the studied variables with serum adiponectin in obese patients have been summarized in table 1. First, baseline levels of adiponectin in obese subjects was compared with that of normal weight subjects, and findings from the independent statistical T test showed lower levels of adiponectin in obese subjects than subjects with normal weight (5.44 ± 1.23 in the obese group vs. 7.56 ± 1.32 in the normal group, $p = 0.021$). These findings indicate decrease in adiponectin levels in obese subjects. The findings also showed a significant negative correlation between serum adiponectin levels and total cholesterol ($p = 0.041$). Significant inverse relationship was also observed between adiponectin and fasting LDL levels ($p = 0.025$). However, the relationship of adiponectin with other lipid profile indicators such as HDL and triglyceride was not significant. On the other hand, the findings showed that decrease in serum adiponectin levels in studied obese subjects is associated with significant increase of fasting glucose ($p = 0.042$). Also decrease in adiponectin concentration was associated with increase of Systolic blood pressure ($p = 0.001$) and diastolic blood pressure ($p = 0.023$) of the studied subjects. Findings from regression analysis showed that although the relationship between adiponectin and each one of the anthropometrical indices is linear and reversed or in other words, an increase in the indicators such as Abdominal circumference, body mass index and Waist to hip circumference ratio (WHO) is associated with decrease in serum adiponectin levels, but the interaction between this hormone and each one of the mentioned variables is not significant separately, which may be due to the high standard deviation or few number of studied samples. Statistical findings also showed that serum adiponectin has a significant inverse relationship with triglyceride ratio to HDL ($p = 0.027$).

Table 1: Anthropometrical characteristics, biochemical variables and a significance levels of each one of the variables with baseline serum adiponectin in the studied obese patients				
intensity correlation (R)	significance level with adiponectin	Range	Standard deviation ± Mean	Variable
R = 0.452	0.005	37-48	43 ± 6	Age (years)
R = 0.191	0.092	84-112	101 ±13.64	Weight (kg)
R = 0.211	0.453	167-185	174 ± 12	Height (cm)
R = 0.573	0.001	11-14	12.83 ± 0.99	Systolic blood pressure (mmHg)
R = 0.449	0.023	7-10	0.96±8.92	Diastolic blood pressure (mmHg)
R = 0.193	0.092	101-132	9.89±106	Abdominal circumference
R = 0.237	0.103	102-121	8.46±108	hip circumference
R = 0.298	0.127	0.96-1.02	0.02±0.99	WHO
R = 0.238	0.210	30-36	3.59±33.83	(kg/m²)Body mass index
R = 0.214	0.113	28-36	4±31.64	Body fat Percentage (%)
R = 0.263	0.221	13-17	2.45±13.91	Visceral obesity
R = 0.497	0.041	118-230	27±182	(mg/dl)Total cholesterol
R = 0.271	0.201	110-240	35±165	(mg/dl) Triglyceride
R = 0.563	0.205	101-165	21±125	Low density lipoprotein(mg/dl)
R = 0.262	0.093	38-48	3.94±44	high density lipoprotein(mg/dl)
R = 0.542	0.042	85-114	9±102	Glucose (mg /dl)
R = 0.534	0.031	2.23-5.43	1.35±3.75	TG/HDL
----	---	4.71-6.30	1.23±5.44	Adiponectin (µg /ml)

4. Discussion / Conclusion

In this study, the role of anthropometrical factors and blood lipid levels on serum Concentration of adiponectin was studied in obese adults. The Initial findings of this study were baseline levels of serum adiponectin being low in obese male adults compared with their peer men with normal weight. This finding indicates the independent impact of obesity on this peptide hormone.

Although this study results support the reduction of adiponectin levels in obese subjects, once the independent relation of each one of factors affecting obesity such as BMI or abdominal circumference was examined with serum adiponectin separately, it became clear that although their relationship with adiponectin was linear and reverse or in other words, increase of the levels of anthropometrical indices is associated with decrease in serum adiponectin levels, this relationship was not statistically significant. These findings show that combination of a set of determinants of obesity with one another leads to reduction of adiponectin levels in obese subjects. In fact, relying on the serum adiponectin levels being lower in obese individuals than subjects with normal weight in this study, it can be concluded that obesity has been a decisive factor in adiponectin reduction in the studied subjects. But since the relationship of adiponectin with each anthropometrical component of body is reverse but non-significant, it may be concluded that although in this study increase of each obesity indicating factor is not determining the reduction of serum adiponectin levels in obese individuals in itself, increase of these obesity determinants levels next to each other reduces serum adiponectin levels in obese subjects significantly compared to subjects with normal weight.

In other studies, reduction of serum or plasma adiponectin levels has been also observed in obese populations (7). Another major aim of this study is to determine the relationship between lipid profile indices and adiponectin levels in the studied subjects. In this regard, the study results showed that despite the inverse linear relationship between serum adiponectin and fasting triglyceride levels, this relationship is not significant. Absence of significant relationship of serum adiponectin with HDL levels was also observed in this study. These finding state that determination of triglycerides levels and blood HDL are not appropriate predictive indicators for serum adiponectin. On the other hand, despite the inverse relationship of serum adiponectin with TG and HDL in this study, non-significance of this relationship may be because of low number of studied samples. But despite the lack of significant relationship of serum adiponectin with HDL and triglyceride levels in these individuals, the statistical tests indicated a significant inverse relationship between total cholesterol and LDL levels and serum adiponectin concentrations. So that increase in blood

total cholesterol level leads to reduction of serum adiponectin as one of the indices of determination of blood cycle lipid levels. In this field, Rubin et al. study showed that increase of total cholesterol levels is associated with decrease in adiponectin levels (8). Despite the lack of significant relationship of adiponectin with triglycerides and HDL indices in this study, statistical findings showed that adiponectin levels of studied obese subjects have a reverse significant relationship with triglycerides ratio to high density lipoprotein (TG / HDL) which is of the determinants of cardiovascular disease (9). This study also showed that age is also reversely related with adiponectin levels so that blood adiponectin concentration would be reduced in line with age increasing. The findings of this study also showed that decrease in serum adiponectin levels is associated with increase of blood pressure and fasting glucose level of the studied subjects. The Inverse relationship of adiponectin with blood pressure in obese patients has also been observed in the study of Lee et al (10).

In an overall summing up, the findings of this study along with most previous findings indicate decrease baseline adiponectin levels in obese subjects compared with those with normal weight. At the end, it can be said that although in the present study on obese subjects, cause for obesity and increase of total cholesterol levels and LDL was associated with significant decrease in adiponectin levels and findings indicate significant inverse correlation of adiponectin and with lipid profile indicators, the relationship of peptide hormone with other components, lipid profile or anthropometrical indices of studied subjects was weak and despite the present findings, there are still unclear points in this regard there which reminds the requirement for further studies with more samples.

5. Acknowledgments

The authors of the article appreciate the research deputy of Azad University of Saveh, Danesh Hematology laboratory of the city as well as all participants in the present study which is the result of the research proposal sponsored by Azad University of Saveh.

6. References

- [1] Kahn CR. Triglycerides and toggling the tummy. *Nat. Genet.* 2000; 25:6–7.
- [2] Steppan CM. The hormone resistin links obesity to diabetes. *Nature.* 2001; 409:307–312.
- [3] Yukihiro Y, Hiroshi H, Ikuo S, Motowo T, Matsuo T, Koichi M et al. Correlation of the adipocyte-derived protein adiponectin with insulin resistance index and serum high-density lipoprotein-cholesterol, independent of body mass index, in the Japanese population. *Clinical Science.* 2002; 103: 137–142.
- [4] Snehalatha C, Yamuna A, Ramachandran A. Plasma Adiponectin Does Not Correlate With Insulin Resistance and Cardiometabolic Variables in Nondiabetic Asian Indian Teenagers. *Diabetes Care.* 2008; 31(12):2374-9.
- [5] Ram W, Sylvie D, Aida G, Kitt P, James D, Sara ET et al. Low Adiponectin Levels in Adolescent Obesity: A Marker of Increased Intramyocellular Lipid Accumulation. *J Clin Endocrinol Metab.* 2003, 88(5):2014–2018.
- [6] Co[^]te M, Maurie[^]ge P, Bergeron J, Alme[^]ras N, Tremblay A, Lemieux I, et al. Adiponectinemia in visceral obesity: impact on glucose tolerance and plasma lipoprotein-lipid levels in men. *J Clin Endocrinol Metab.* 2005; 90:1434–9.
- [7] Cnop M, Havel PJ, Utzschneider KM, Carr DB, Sinha MK, Boyko EJ et al. Relationship of adiponectin to body fat distribution, insulin sensitivity and plasma lipoproteins: evidence for independent roles of age and sex. *Diabetologia.* 2003; 46: 459–469.
- [8] Rubin DA, McMurray RG, Harrell JS, Hackney AC, Thorpe DE, Haqq AM. The association between insulin resistance and cytokines in adolescents: the role of weight status and exercise. *Metabolism.* 2008; 57(5):683-690.
- [9] Jiang R, Schulze MB, Li T, Rifai N, Stampfer MJ, Rimm EB: Non-HDL cholesterol and apolipoprotein B predict cardiovascular disease events among men with type 2 diabetes. *Diabetes Care.* 2004; 27:1991-1997.
- [10] Li JQ, Li MY, Dai YX, Liu YM, Zheng J, Liu WB. Serum adiponectin and free fatty acid profile in essential hypertensive patients with or without metabolic syndrome. *Zhonghua Xin Xue Guan Bing Za Zhi.* 2008; 36(3):232-5.