

## Effects of Nitrogen Fertilizer on Wild Oat (*Avena fatua*) Competition with Wheat (*Triticum aestivum*)

Jafar Pourreza<sup>1</sup> and Abdollah Bahrani<sup>2</sup>

<sup>1,2</sup>Department of Agriculture, Ramhormoz Branch, Islamic Azad University, Ramhormoz, Iran

**Abstract.** A field experiment was conducted to quantify the effect of nitrogen fertilizer on wheat and wild oat competition. The experiment was designed as a randomized complete block with a split plot arrangement with nitrogen fertilizer as main plot and wild oat density as subplot. The relationship between wheat yield loss percentage versus wild oat density was described using a rectangular hyperbola model. Initial slope ( $i$ ) of the rectangular hyperbola model was significantly greater when nitrogen fertilizer was applied. Moreover, for the rectangular hyperbola model, there was significant effect of nitrogen application on estimated maximum wheat yield loss ( $A$ ). The application of 150 N Kg ha<sup>-1</sup> (N<sub>1</sub> treatment) before crop seeding resulted in a greater higher competitive ability of wild oat than other treatments. The slope ( $b$ ) of the linear model representing the relationship between wild oat density and relative wild oat biomass significantly greater when nitrogen fertilizer applied.

**Keywords:** Wild oat, nitrogen, wheat, competition.

### 1. Introduction

Generally weeds are always considered harmful plants. Weeds are one of the biggest threats to agriculture [1]-[3]. Wild oat was described as a vigorously growing weed with a capability to attain greater height, and establish and develop extensive leaf area and horizontal branches when moisture and nutrients are not limiting [4]-[7]. These morphological and physiological characteristics of wild oat allow it to shade and suppress the growth of its neighbours to a level that causes yield reduction [8].

Fertilization is an important agronomic strategy used extensively to increase crop yield. Nevertheless, although nutrients clearly promote crop growth, many studies have shown that, in some cases, fertilizers benefit weeds more than crops [9]. For example, Carlson and Hill [10] found that the addition of N fertilizer to wild oat-infested wheat increased the density of wild oat panicles without increasing crop yield. However, Satorre and Snaydon [11] showed that N fertilizer reduced the severity of competition experienced by wild oat from six spring cereals. The increase in weed competition at higher N rates has been suggested to be related to an increase in the efficiency of nutrient accumulation and use by weeds [9]. In addition, Brandt et al. [12] reported differences in weed response to soil fertility for wheat and canola (*Brassica napus* L.), suggesting that this may be exploited through the development of agronomic systems that stimulate crop growth over weed growth. Moreover, Knezevic et al. [13] noted that insufficient N can reduce corn (*Zea mays* L.) tolerance to weeds and can lengthen the critical period for weed control. Fertilizer application can have an influence on the composition of the weed community. In this sense, Jornsgard et al. [14] demonstrated an interaction between the growth of individual weed species and the level of N fertilizer in cereal crops. They found that common lambsquarters (*Chenopodium album* L.) in competition with spring barley has a lower N optimum than does the crop, whereas burning nettle (*Urtica urens* L.) has a higher N optimum than does the crop.

The objective of this was to investigate the effect of nitrogen fertilization application in simulating wheat (*Triticum aestivum*) yield loss caused by wild oat (*Avena fatua*) interference.

## 2. Materials and Methods

### 2.1. Study site, Experimental Design, Treatments and crop operations

A field experiment was conducted at agricultural research station of Islamic Azad University, Ramhormoz branch (latitude 31.16 °N, longitude 49.36 °E and 151 m asl), south-western of Iran. Planting date was 6 December in 2008. The site has a silty clay soil with a mean annual precipitation 320 mm, mean solar radiation of 19.1 MJ m<sup>-2</sup> d<sup>-1</sup>, mean maximum temperature of 32.7 °C and mean minimum temperature of 19.5 °C. The climate is warm and dry. The experiment was designed as a randomized complete block with a split plot arrangement with nitrogen fertilizer ( N<sub>0</sub>: unfertilized control, N<sub>1</sub>: the use of 150 N Kg ha<sup>-1</sup> before wheat seeding and N<sub>2</sub>: the use of 50 N Kg ha<sup>-1</sup> before wheat seeding + 100 N Kg ha<sup>-1</sup> at late tillering stage of wheat) as main plot and wild oat density (D<sub>0</sub>: weed- free control, D<sub>1</sub>: 25, D<sub>2</sub>: 50, D<sub>3</sub>: 75 and D<sub>4</sub>: 100 plants m<sup>-2</sup>) as subplot with three replications. Seeding rates for wild oat were based on target stand densities, 1,000 seed weight and assumption of 10% recruitment. The wild oat seeds was sown at a depth of approximately 2 to 3 cm, between 2 planting row of wheat. After wild oat seed was sown (but before wheat seeding) granular urea fertilizer (46-0-0) was broadcast mechanically onto the main plot nitrogen treatments. Residual rate of nitrogen fertilizer (100 Kg ha<sup>-1</sup>) which related to N<sub>2</sub> treatment was applied late tiller stage of wheat. An unfertilized control was also included. Subplot size was 6×1.25 m and all subplots were separately by a 2.5 m wide alley. Individual plots consisted of six rows, 6 m long and spaced 25 cm apart. All subplots were sown by a seed drill at a wheat population density of 400 seed m<sup>-2</sup>. In general, weed control throughout the growing season was done by hand weeding. Insects were controlled by the appropriate chemicals. All the experiments were conducted under well-watered conditions. The plots were irrigated after 60-mm cumulative pan (Class A) evaporation and irrigation amount was based on soil moisture depletion. Therefore, there was no effect of flooding or water deficit stresses.

### 2.2. Measurements

In late January, when both crop and weeds had completed emergence, an area of 1 m<sup>2</sup> was marked in the centre of each plot. Weed seedling were counted and thinned as necessary to achieve desired densities. The six more samples (by harvesting 0.25 m<sup>2</sup>) were taken throughout growing season for determination of above ground dry weight(drying oven at 70 °C for 48 h) of both wheat and wild oat. In late May wheat and wild oat plants from the previously marked areas were harvested by hand, then total dry matter and grain yield of wheat and total above ground weight of wild oat determined.

### 2.3. Statistical analysis

Percent wheat yield loss was calculated using the following formula;

$$YL = [(Y_{wf} - Y_{weedy})/Y_{wf}] \cdot 100 \quad (1)$$

where YL is percent yield loss, Y<sub>wf</sub> is weed-free yield, and Y<sub>weedy</sub> is yield in weed (wild oat)-infested plots. Percent relative wild oat biomass was calculated using the following formula [3]:

$$RelWb = [W_{wo}/(W_{wo} + W_w)] \cdot 100 \quad (2)$$

where RelWb is relative wild oat biomass, W<sub>wo</sub> is aboveground dry biomass of wild oat, and W<sub>w</sub> is aboveground dry biomass of wheat in the same plot. Data were subjected to ANOVA using the PROC GLM procedure in SAS [15].

For percent wheat yield loss and percent relative wild oat biomass, on the basis of the results obtained by an ANOVA, data sets were subjected to regression analysis using actual wild oat density as the independent variable. A rectangular hyperbola model was fitted where possible [16]:

$$YL = i d/[1 + (i d/A)] \quad (3)$$

where YL is percent wheat yield loss, *d* is wild oat density (plants. m<sup>-2</sup>), *i* is the initial slope, and *A* is the asymptote. The model was fitted to the data using the PROC NLIN procedure of SAS [15].

For percent relative wild oat biomass, A linear model was used as follows:

$$RrLW_b = b \times d \quad (4)$$

where *d* is wild oat density (plants. m<sup>-2</sup>) and *b* is the slope of linear model. The data for each parameter were individually subjected to ANOVA and the significant means were separately by the LSD test.

### 3. Results and Discussion

The application of nitrogen fertilizer increased the competitive effect of wild oat on spring wheat. The initial slope ( $i$ ) of the rectangular hyperbola model was significantly greater when nitrogen fertilizer was applied (Table 1). Moreover, for the rectangular hyperbola model, there was significant effect of nitrogen application on estimated maximum wheat yield loss ( $A$ ). Carlson and Hill [10] and Ross and Van Acker [3] also found that wheat yield loss due to wild oat interference was greater when nitrogen fertilizer was applied. However, The application of 150 N Kg ha<sup>-1</sup> (N<sub>1</sub> treatment) before crop seeding resulted in a greater higher competitive ability of wild oat than other treatments (Table 1).

Table 1: Parameter estimates for rectangular hyperbola fitted to data of wheat yield loss as a function of wild oat density and as affected by nitrogen fertilizer.

	$i \pm se^a$	$A \pm se$	RMSE <sup>b</sup>	R <sup>2</sup>
N <sub>0</sub>	0.82 ± 0.20	40.48 ± 7.44	1.89	0.99
N <sub>1</sub>	1.27 ± 0.03	69.57 ± 1.33	0.30	0.99
N <sub>2</sub>	1.08 ± 0.18	61.85 ± 1.33	1.94	0.99
<sup>a</sup> Standard error of parameter estimates <sup>b</sup> Root Mean Square Error				

The comparison of initial slope ( $i$ ) and estimated maximum wheat yield loss ( $A$ ) between nitrogen treatments indicated that application of nitrogen than unfertilized control (N<sub>0</sub> treatment) increased the rates of these coefficients. However, the coefficient rates were greater in N<sub>1</sub> treatment (the use of 150 N Kg ha<sup>-1</sup> before crop seeding) than N<sub>2</sub> (the use of 50 N Kg ha<sup>-1</sup> before crop seeding + 100 N Kg ha<sup>-1</sup> at late tillering stage of wheat). This result suggests more nitrogen application before crop seeding increased the wheat yield loss than that of treatments, which would be due to the more germinability of weed seeds [5]. Some studies showed that nitrogen fertilizer can stimulate the weed seed dormancy (2, 3), so because of more weed plant in area, wheat yield affected and reduced.

This result is of practical concern for farmers in this region because wild oat is a primary pest, and extra use of nitrogen regardless of managed recommendations can benefit weed growth.

Table 2: Parameter estimates for linear models fitted to data of wild oat relative biomass as a function of wild oat density and affected by nitrogen fertilizer.

	$b \pm se^a$	RMSE <sup>b</sup>	R <sup>2</sup>
N <sub>0</sub>	0.13 ± 0.01	1.57	0.97
N <sub>1</sub>	0.19 ± 0.02	2.54	0.96
N <sub>2</sub>	0.17 ± 0.01	2.00	0.97
<sup>a</sup> Standard error of parameter estimates <sup>b</sup> Root Mean Square Error			

The application of nitrogen fertilizer (N<sub>1</sub> and N<sub>2</sub> treatments) resulted in a greater wild oat biomass accumulation relative to wheat biomass than an unfertilized control treatment (table 2). The slope ( $b$ ) of the linear model representing the relationship between wild oat density and relative wild oat biomass significantly greater when nitrogen fertilizer applied. This result would help explain why the effect of wild oat on wheat yield increased when nitrogen fertilizer was applied. It also suggests that wild oat more effectively exploited surface-applied nitrogen fertilizer than did spring wheat. Di Tomaso [9] observed that broadcast nitrogen fertilizer applied under high weed densities tended to increase weed growth with little benefit to crops. Similarly, Blackshaw et al. [2], [17]) and Ross and Van Acker [3] have shown that nitrogen fertilizer placement (relative to the crop) can make spring wheat more competitive against weeds. In this study, wild oat seedlings may have had preferential access to surface-applied nitrogen because the wild oat seed was broadcast and incorporated nearer the soil surface (2 to 3 cm deep) than the wheat (4–6 cm). As well, Pavlychenko and Harrington [18] observed markedly reduced rooting systems in wheat vs. wild oat under competition. They found average total root length of wheat and wild oat to be 48.6 and 252.3 m, respectively, when the two species were grown together, while the average root length for wheat grown

alone was 90 m. The more extensive root system of wild oat may allow it to continue to preferentially access and exploit soil nitrogen through the growing season. This may be true for other nutrients as well. For example, Callow et al. [19] found that when KCl fertilizer was applied to field plots, wild oat was more competitive with spring wheat.

Blackshaw et al. [2] and Sexsmith and Pittman [6] found that nitrogen fertilizer significantly increased wild oat biomass in spring wheat, and this translated into significantly greater wild oat seed return.

Wheat grain yield was significantly affected by wild oat densities, nitrogen levels and by their interactions (table 3 ). So, main effects of treatments and wild oat density  $\times$  nitrogen interaction means are presented (table 4 and 5). The data in table 4 show that for the nitrogen levels, the maximum (302.4 g m<sup>-2</sup>) grain yield was observed in N<sub>2</sub> treatment while lowest of that related to unfertilized control (N<sub>0</sub>). Among the wild oat densities the highest (332.7 g m<sup>-2</sup>) grain yield was recorded in weed-free treatment (D<sub>0</sub>), while lowest (203.9 g m<sup>-2</sup>) grain yield was recorded in 100 wild oat seed m<sup>-2</sup> treatment (D<sub>4</sub>).

Table 3: Analysis of variance of wheat grain yield and total dry matter of wild oat data as affected by different nitrogen levels and wild oat densities.

Source	DF	Mean square	
		Wheat grain yield	Total dry matter of wild oat
Block	2	2.95 <sup>NS</sup>	45.26 <sup>NS</sup>
Nitrogen	2	30800.25 <sup>**</sup>	3366.86 <sup>**</sup>
Error	4	22.75	15.63
Wild oat density	4	24936.30 <sup>**</sup>	35564.85 <sup>*</sup>
Nitrogen $\times$ wild oat density	8	597.48 <sup>**</sup>	402.42 <sup>**</sup>
Error	24	20.24	8.64
CV %		1.79	2.88

\*Difference significant (p<0.05); \*\* difference significant (p<0.01); NS, not significant

Table 4: The wheat grain yield (g m<sup>-2</sup>) as affected by different nitrogen levels and wild oat densities.

N	Wild oat densities m <sup>-2</sup>					N means <sup>aa</sup>
	0	25	50	75	100	
N <sub>0</sub>	281.3 <sup>d</sup>	238.0 <sup>f</sup>	200.0 <sup>h</sup>	190.0 <sup>h</sup>	193.3 <sup>de</sup>	220.5 <sup>c</sup>
N <sub>1</sub>	316.6 <sup>b</sup>	246.6 <sup>e</sup>	212.3 <sup>g</sup>	190.0 <sup>h</sup>	173.3 <sup>i</sup>	227.8 <sup>b</sup>
N <sub>2</sub>	400.3 <sup>a</sup>	320.6 <sup>b</sup>	294.3 <sup>c</sup>	251.6 <sup>e</sup>	245.0 <sup>e</sup>	302.4 <sup>a</sup>
Densities means <sup>bb</sup>	332.7 <sup>a</sup>	268.4 <sup>b</sup>	235.5 <sup>c</sup>	210.5 <sup>d</sup>	203.9 <sup>e</sup>	

<sup>aa</sup> LSD(0.05) for N means = 4.50  
<sup>bb</sup> LSD(0.05) for densities means = 4.37

Table 5. Total dry matter (g m<sup>-2</sup>) of wild oat as affected by different nitrogen levels and wild oat densities.

N	Wild oat densities m <sup>-2</sup>				N means <sup>aa</sup>
	25	50	75	100	
N <sub>0</sub>	76.3 <sup>i</sup>	95.3 <sup>g</sup>	121.0 <sup>d</sup>	126.0 <sup>a</sup>	83.8 <sup>c</sup>
N <sub>1</sub>	102.3 <sup>f</sup>	122.6 <sup>d</sup>	155.3 <sup>b</sup>	181.3 <sup>a</sup>	106.2 <sup>b</sup>
N <sub>2</sub>	90.3 <sup>h</sup>	114.0 <sup>e</sup>	147.3 <sup>c</sup>	179.3 <sup>a</sup>	112.3 <sup>a</sup>
		110.6 <sup>c</sup>	141.4 <sup>b</sup>	162.2 <sup>a</sup>	
Densities means <sup>bb</sup>	89.6 <sup>d</sup>	110.6 <sup>c</sup>	141.4 <sup>b</sup>	162.2 <sup>a</sup>	

<sup>aa</sup> LSD(0.05) for N means = 2.40  
<sup>bb</sup> LSD(0.05) for densities means = 2.83

For the interaction of nitrogen with the wild oat densities, the highest grain yield (400.3 g m<sup>-2</sup>) was observed in N<sub>2</sub> (use of 50 N Kg ha<sup>-1</sup> before crop seeding + 100 N Kg ha<sup>-1</sup> at late tillering stage of wheat ) treatment  $\times$  0 wild oat seed. The minimum grain yield (173.3 g m<sup>-2</sup>) was observed in N<sub>1</sub> ( the use of 150 N

Kg ha<sup>-1</sup> before crop seeding) × 100 wild oat seed m<sup>-2</sup>. The mean weed-free wheat yield at N<sub>0</sub> was lower than N<sub>1</sub> and N<sub>2</sub> treatments (Table 4). Weed-free wheat yields seem related to pre-seeding and post-seeding soil nitrogen levels, which were greatest for N<sub>2</sub> that nitrogen application in soil was applied at the two stages. The higher wild oat densities have had wheat yield loss than that of lower wild oat densities. Grain yield of wheat (with or without weed competition) was higher at N<sub>2</sub> treatment compared with N<sub>1</sub> treatment (table 4).

Study results support previous findings indicating that many agricultural weed species are as responsive as or more responsive than crops to higher soil N levels [2, 20]. Thus, indiscriminate N fertilizer use has the potential to benefit weeds at the expense of crops [3]. Grower in south western of Iran sometimes apply N fertilizer in the previous fall before seeding spring wheat as a means of decreasing the workload at planting. Results indicate that this is not necessarily a bad practice in terms of wheat yield in absence of weeds. However, wheat infested with competitive weed species sometimes had lower yield with preplant (N<sub>1</sub>) than with multistage-applied N (N<sub>2</sub>). Multistage- applied N compared with preplant –applied N was never worse in terms of weed density, weed biomass (Table 5), wheat yield (table 5), wheat yield loss (table 2) and relative wild oat biomass loss (table 3), indicating that it is a less-risky N application timing in terms of both weed management wheat yield response.

The nitrogen application (N<sub>1</sub> and N<sub>2</sub> treatments) compared with unfertilized control treatment (N<sub>0</sub>) tended to have larger and more consistent effect on weed growth and crop weed competition. Although, the wheat yield loss was increased by N application, but a such reduction was more recorded at once N application than multistage- applied N. The findings reported by Pedreros [21] and Blackshaw et al. [2] support our conclusions.

Our results are strongly supported by other researchers [2], [3], [5] who indicated application of nitrogen can be effective factor to effect the competitive ability weed and crop. In contrast, other studies found that N application method [22] or dose [23] had little effect on crop-weed competition. In additional, results may be crop and weed specific [2]. Common Chickweed (*Stellaria media* L.) interference with potato (*Solanum tuberosum*) was reduced with higher N levels, but the opposite result occurred when it was competing with wheat. Journsgard et al [14] report that weed biomass in barely (*Hordeum vulgare* L.) and winter wheat could be increased, unchanged or reduced with increased soil N, depending on the weed and crop.

## 4. Conclusions

The results of this study suggest that the application of nitrogen fertilizer to wild oat-infested wheat fields increases wild oat competitions. Although, the wheat yield loss was increased by N application, but a such reduction was more recorded at once N application than multistage- applied N. Greater knowledge of the effect of N on weeds and crop grown in competitive mixtures may allow a better understanding of why differences occurred among previous studies and would aid development of fertilization strategies to reduce weed competition with crops. Manipulation of crop fertilization not only had potential to protect crop yield but also may contribute to long term reduction in weed populations. N fertilizer is known to break the dormancy of certain weed species [9] and thus may directly affected weed infestation densities.

## 5. References

- [1] El-Khatib, A. A. and A. K. Hegazy. 1999. Growth and physiological responses of wild oats to the allelopathic potential of wheat. *Acta Agronomica Hungarica* 47: 11-18.
- [2] Blackshaw, R. E., L. J. Molnar, and H. H. Janzen. 2004. Nitrogen fertilizer timing and application method affect weed growth and competition with spring wheat. *Weed Sci.* 52:614–622.
- [3] Ross, D. M., and R. C. Van Acker. 2005. Effect of nitrogen fertilizer and landscape position on wild oat (*Avena fatua*) interference in spring wheat. *Weed Science*, 53:869–876
- [4] Heyne, E. G. 1987. *Wheat and Wheat Improvement*, 2nd edition Madison, Wisconsin, USA.
- [5] Dhima, K.V., Eleftherohorinos, I.G., and Vasilakoglou, I.B. 2000. Interference between *Avena sterilis*, *Phalaris minor* and five barley cultivars. *Weed Res.* 40:549-559.
- [6] Sexsmith, J. J. and U. J. Pittman. 1962. Effect of nitrogen fertilizers on germination and stand of wild oats. *Weeds* 11:99–101.

- [7] Dhima, K. V. and I. G. Eleftherohorinos. 2001. Influence of nitrogen on competition between winter cereals and sterile oat. *Weed Sci.* 49:77– 82.
- [8] Cudney, D. W., L. S. Jordan and A E. Hall. 1991. Effect of wild oat (*Avena fatua*) infestations on light interception and growth rate of wheat (*Triticum aestivum*). *Weed Sci.* 39: 175-179.
- [9] DiTomaso, J. M. 1995. Approaches for improving crop competitiveness through the manipulation of fertilization strategies. *Weed Sci.* 43:491– 497.
- [10] Carlson, H. L. and J. E. Hill. 1986. Wild oat (*Avena fatua*) competition with spring wheat: effects of nitrogen fertilization. *Weed Sci.* 34:29– 33.
- [11] Satorre, E. H. and R. W. Snaydon. 1992. A comparison of root and shoot competition between spring cereals and *Avena fatua* L. *Weed Res.* 32: 45–55.
- [12] Brandt, R., H. H. Janzen, and R. E. Blackshaw. 2000. Ranking weed response to added nitrogen and phosphorous. Abstracts of the III International Weed Science Congress; Foz do Iguassu, Brazil; 6–11 June, 2000. 26 p.
- [13] Knezevic, S., S. Evans, C. Shapiro, and J. Lindquist. 2000. Effect of nitrogen on critical period of weed control in corn. Abstracts of the III International Weed Science Congress; 6–11 June, 2000; Foz do Iguassu, Brazil. 52p.
- [14] Jornsgard, B., K. Rasmussen, J. Hill, and J. L. Christiansen. 1996. Influence of nitrogen on competition between cereals and their natural weed populations. *Weed Res.* 36:461–470.
- [15] SAS Institute Inc., 1989. SAS user' guide: Statics, Version 6, 4<sup>th</sup> editions, SAS Inst., Inc., Cary, N.C.
- [16] Cousens, R. 1985. A simple model relating yield loss to weed density. *Ann. Appl. Biol.* 107:239–252.
- [17] Blackshaw, R. E., G. Semach, and H. H. Janzen. 2002. Fertilizer application method affects nitrogen uptake in weeds and wheat. *Weed Sci.* 50:634–641.
- [18] Pavlychenko, T. K. and J. B. Harrington. 1935. Root development of weeds and crops in competition under dry farming. *Sci. Agric.* 16:151–160.
- [19] Callow, K. A., D. A. Derksen, C. A. Grant, and R. C. Van Acker. 1999. The impact of monoammonium phosphate and potassium chloride on wild oat (*Avena fatua* L.) competition in zero-till spring wheat and flax. Pages 57–59 in 21st Annual Manitoba-North Dakota Zero Tillage Workshop, Brandon, MB, Canada. Brandon, MB, Canada: Manitoba- North Dakota Zero-tillage Association.
- [20] Supasilapa, S., B. T. Steer, and S. P. Milroy. 1992. Competition between lupin (*Lupinus angustifolia* L.) and great brome (*Bromus diandrus* Roth.): development of leaf area, light interception and yields. *Aust. J. Exp. Agric.* 32:71–81.
- [21] Pedreros, L.A. 2001. Wild oat (*Avena fatua* L.) and Italian ryegrass (*Lolium multiflorum* Lam.) effect on wheat yield at two locations. *Agricultura Tecnica.* 61: 294-305.
- [22] Cochran, V. L., L. A. Morrow, and R. D. Schirman. 1990. The effect of N placement on grass weeds and winter wheat in three tillage systems. *Soil Tillage Res.* 18:347–355.
- [23] Gonzalez Ponce, R. 1998. Competition between barley and *Lolium rigidum* for nitrate. *Weed Res.* 38:453–460.