

Study on Specie Diversity, Zoogeographical Distribution and Ecological Properties of the Miridae (Hemiptera) Family in the Hulun Buir City, Inner Mongolia of China

Kai Shi¹⁺, Yuanyuan Li¹ and Changhua Bao²

¹ Agricultural College, Inner Mongolia University for the Nationalities, Tongliao, Inner Mongolia, 028043, China

² Medical College, Inner Mongolia University for the Nationalities, Tongliao, Inner Mongolia, 028043, China

Abstract. The main aim of this study was to understand the species diversity composition, zoogeographic distribution and ecological properties of the family Miridae (Hemiptera) from the Hulun Buir city. Our results showed there were 51 genera 122 species, including three new record species of China and one endemic species of China. In the world animal geographical fauna, the composition of Palaearctic region was dominant. There were 79 species only distributed there, about 64.75 % of the total known. Studies of ecological properties had shown the most species were distributed in 48 ° 01' -49 ° N and the most appropriate altitudinal range was 601-800 m.

Keywords: Miridae, species diversity, zoogeographical distribution, ecological properties.

1. Introduction

The Hulun Buir city is located in 47° 05' - 53° 20' N and 115° 31' - 126° 04' E, Northeast of Inner Mongolia of China; and having an area of approximately 253,000 square kilometers[1]. Its territory has famous Hulun Buir Grassland. The city is a transitional zone from the forest meadows of Northeast of China to arid grassland. In recent years, due to the large area of farmland reclamation and mine development, coupled with the rapid development of tourism, the Hulun Buir city showing habitat fragmentation, has become Chinese typical ecological fragile area.

Miridae is the first largest family of Heteroptera of Hemiptera. Up to now, more than 1,400 genera, 11,139 species have been described worldwide (Schuh, 2002-2013, <http://research.amnh.org/pbi/catalog/>). Most Miridae are herbivorous, especially *Lygus* spp., *Trigonotylus* spp. and *Apolygus* spp. likely become threatening pasture pests. Our study focuses on the following three aspects, studying the species diversity composition, zoogeographic distribution and ecological properties of Miridae in order to provide some supporting data for further the relevant biogeography, biodiversity conservation, ecology research etc.

2. Materials and Methods

Our data were compiled from three resources: from all the literatures published up to now; from all the collected records between June and September of 2011-2015 deposited in Inner Mongolia University for the Nationalities; from accumulation label records of specimens and/or types deposited in the Nonnaizab Entomology Research Center of Inner Mongolia Normal University. According to the literatures [2]-[4], all the specimens were sorting, classification, verification and identification. In addition, we also conducted the

⁺ Corresponding author. Tel.: +86-475-8314809
E-mail address: nmshikai@126.com

zoogeographical distribution analysis according to classical six zoogeographical regions [5]. Using Excel 2010 statistical software, we made ecological properties analysis to Miridae of Hulun Buir.

Table 1: Miridae species composition of Hulun Buir city and their zoogeographical distribution

PA: Palaearctic region, OR: Oriental region, NEA: Nearctic region, ET: Ethiopian region, ● New record species of China; ▲ endemic species of China

Subfamily	Genus	Species	Zoogeographical Distribution	
Deraeocorinae	<i>Deraeocoris</i>	<i>D. annulipes</i> (Herrich-Schaeffer)	PA	
		<i>D. ater</i> (Jalovlev)	PA	
		<i>D. kerzhneri</i> Josifov	PA	
		<i>D. morio</i> (Boheman)	PA	
		<i>D. olivaceus</i> (Fabricius)	PA	
		<i>D. pallidicornis</i> Josifov	PA	
		<i>D. punctulatus</i> (Fallén)	PA, NEA	
		<i>D. salicis</i> Josifov	PA	
		<i>D. scutellaris</i> (Fabricius)	PA	
		Mirinae	<i>Adelphocoris</i>	<i>A. fasciaticollis</i> Reuter
<i>A. ferrugineus</i> Hsiao	PA			
<i>A. laeviusculus</i> Vinokurov	PA, OR			
<i>A. lineolatus</i> (Goeze)	PA, OR			
<i>A. melanocephalus</i> Reuter	PA			
<i>A. nigritylus</i> Hsiao	PA, OR			
<i>A. obliquefasciatus</i> Lindberg	PA, OR			
<i>A. ponghvariensis</i> Josifov	PA, OR			
<i>A. quadripunctatus</i> (Fabricius)	PA, OR			
<i>A. reicheli</i> (Fieber)	PA			
<i>A. rufescens</i> Hsiao	PA, OR			
<i>A. tenebrosus</i> (Reuter)	PA			
<i>A. triannulatus</i> (Stål)	PA			
<i>Apolygus</i>	<i>A. lucorum</i> (Meyer-Dür)			PA, OR, NEA
	<i>A. nigronasutus</i> (Stål)			PA
	<i>A. nigrovirens</i> (Kerzhner)			PA
	<i>A. spinolae</i> (Meyer-Dür)			PA, OR
	<i>Capsodes</i>			<i>C. gothicus</i> (Linnaeus)
			<i>C. cinctus</i> (Kolenati)	PA, NEA
<i>C. pilifer</i> Remane			PA	
<i>Capsus</i>	<i>C. wagneri</i> Remane		PA	
	<i>Lygidea</i>		<i>L. illota</i> (Stål)	PA
			<i>Lygocoris</i>	<i>L. pabulinus</i> (Linnaeus)
<i>Lygus</i>	<i>L. adpersus</i> (Schilling)			PA
	<i>L. gemellatus</i> (Herrich-Schaeffer)		PA	
	<i>L. orientis</i> Aglyamzyanov		PA, NEA	
	<i>L. pratensis</i> (Linnaeus)		PA	
	<i>L. punctatus</i> (Zetterstedt)		PA, OR, NEA	
<i>Neolygus</i>	<i>L. rugulipennis</i> Poppius		PA	
	<i>L. sibiricus</i> Aglyamzyanov		PA	
	<i>L. wagneri</i> Remane		PA	
	<i>N. chinensis</i> (Lu and Yasunaga)		PA	
	<i>Orthops</i>		<i>O. forelii</i> Fieber ●	PA
			<i>O. mutans</i> (Stål)	PA, OR
			<i>O. scutellatus</i> Uhler	PA
	<i>Phytocoris</i>		<i>P. insignis</i> Reuter	PA
			<i>P. nowickyi</i> Fieber	PA
			<i>Polymerus</i>	<i>P. brevicornis</i> (Reuter)
	<i>P. carpathicus</i> (Horvath)			PA
	<i>P. cognatus</i> (Fieber)			PA
<i>P. funestus</i> (Reuter)	PA			
<i>P. palustris</i> (Reuter)	PA, OR			
<i>P. pekinensis</i> Horvath	PA, OR, NEA			
<i>P. unifasciatus</i> (Fabricius)	PA, OR			
<i>A. signatus</i> Reuter*	PA, NEA			
<i>Leptopterna</i>	<i>L. albescens</i> (Reuter)	PA		
	<i>L. kerzhneri</i> Vinokurov	PA		
	<i>L. xilingolana</i> Jorigtoo and Nonnaizab ▲	PA		
	<i>M. gracilis</i> (Sahlberg)	PA		
<i>Myrmecoris</i>	<i>N. sibirica</i> Golub	PA, OR		
	<i>Stenodema</i>	<i>S. calcarata</i> (Fallén)	PA	
<i>S. holsata</i> (Fabricius)		PA		
<i>S. mongolia</i> Nonnaizab and Jorigtoo		PA		
<i>S. parvulum</i> Zheng		PA, OR		
<i>S. pilosa</i> (Jakovlev)		PA, OR		
<i>S. sericans</i> (Fieber) ●		PA		
<i>S. sibirica</i> Bergroth		PA		
<i>S. trispinosa</i> Reuter		PA, NEA		
<i>S. turanic</i> Reuter		PA		

		<i>S. virens</i> (Linnaeus)	PA, NEA
	<i>Teratocoris</i>	<i>T. saundersi</i> Douglas and Scott	PA, NEA
	<i>Trigonotylus</i>	<i>T. caelestialium</i> (Kirkaldy)	PA, NEA, OR
		<i>T. cremeus</i> Golub	PA
		<i>T. longitarsus</i> Golub	PA
		<i>T. ruficornis</i> (Geoffroy)	PA, NEA, ET
Orthotylinae	<i>Anapus</i>	<i>A. kirschbanumi</i> Stål	PA, NEA
	<i>Euryopicoris</i>	<i>E. nitidus</i> (Meyer-Dür)	PA
	<i>Halticus</i>	<i>H. apterus</i> (Linnaeus)	PA, NEA
		<i>H. pusillus</i> (Herrich-Schaeffer)	PA
	<i>Labops</i>	<i>L. bami</i> Kulik	PA
		<i>L. nivchorum</i> Kerzhner	PA
		<i>L. sahlbergi</i> (Fall )	PA, NEA
	<i>Myrmecophyes</i>	<i>M. alboornatus</i> (St )	PA
	<i>Orthocephalus</i>	<i>O. funestus</i> Jakovlev	PA
	<i>Strongylocoris</i>	<i>S. leucocephalus</i> (Linnaeus)	PA
	<i>Blepharidopterus</i>	<i>B. angulatus</i> (Fall )	PA, NEA
	<i>Cyllecoris</i>	<i>C. equestris</i> St 	PA
	<i>Cyrtorhinus</i>	<i>C. caricus</i> (Fall )	PA, NEA
	<i>Excentricus</i>	<i>E. planicornis</i> (Herrich-Schaeffer)	PA
	<i>Globiceps</i>	<i>G. flavomaculatus</i> (Faricius)	PA
	<i>Labopidea</i>	<i>L. algens</i> (Vinokurov)	PA
	<i>Mecomma</i>	<i>M. dispar</i> (Boheman)	PA
	<i>Orthotylus</i>	<i>O. flavosparsus</i> (Sahlberg)	PA, NEA
		<i>O. interpositus</i> Schmidt	PA
		<i>O. oschanini</i> Reuter	PA
		<i>O. parvulus</i> Reuter	PA
	<i>Ulmocyllus</i>	<i>U. virens</i> Seidenstucker	PA
Phylinae	<i>Hallodopus</i>	<i>H. sibiricus</i> Poppius	PA
	<i>Acrotelus</i>	<i>A. pilosicornis</i> (Reuter)	PA
	<i>Atomoscelis</i>	<i>A. asiatica</i> (Josifov)	PA
		<i>A. onustus</i> (Fieber)	PA, NEA
	<i>Chlamydatus</i>	<i>C. pulicarius</i> (Fall )	PA, NEA
		<i>C. pullus</i> (Reuter)	PA, NEA
	<i>Compsidolon</i>	<i>C. pumilum</i> (Jakovlev)	PA, NEA
	<i>Criocoris</i>	<i>C. crassicornis</i> (Hahn)	PA
		<i>C. quadrimaculatus</i> (Fall )	PA
		<i>C. sibiricus</i> Kerzhner	PA
	<i>Europiella</i>	<i>E. leucopus</i> (Kerzhner)	PA
	<i>Eurycolpus</i>	<i>E. flaveolus</i> (St )	PA
	<i>Excentricoris</i>	<i>E. pictipes</i> (Reuter)	PA
	<i>Macrotylus</i>	<i>M. mundulus</i> (St )	PA
		<i>M. zinovievi</i> Kezhner	PA
	<i>Phaeochiton</i>	<i>P. caraganae</i> (Kerzhner)	PA
	<i>Plagiognathus</i>	<i>P. chrysanthemii</i> (Wolff)	PA, NEA
		<i>P. collaris</i> (Matsumura)	PA
	<i>Psallopsis</i>	<i>P. kirgicus</i> (Becker)	PA
	<i>Psallus</i>	<i>P. betuleti</i> (Fall )	PA, NEA
		<i>P. fallenii</i> Reuter	PA, NEA
		<i>P. ulmi</i> Kerzhner and Josifov	PA
	<i>Sacculifer</i>	<i>S. picticeps</i> Kerzhner	PA
	<i>Pilophorus</i>	<i>P. cinnamopterus</i> (Kirschbaum)	PA, NEA
		<i>P. clavatus</i> (Linnaeus)	PA, NEA
		<i>P. setulosus</i> Horvath	PA

3. Results

3.1. Species diversity composition and zoogeographical distribution of Miridae

A total of 122 species from 4 subfamilies 51 genera were recorded from Hulun Buir city. Among of them, *Orthops forelii* Fieber, *Anapus kirschbanumi* St  and *Stenodema sericans* Fieber were new record species of China; and *Leptopterna xilingolana* Jorigtoo and Nonnaizab was the endemic species of China. The subfamily Mirinae had the most abundant species diversity (65 species), followed by Phylinae (26 species). The genus *Adelphocoris* Reuter had the most species (13 species).

The analysis of zoogeographical distribution showed all the genera and species were distributed in the Palaearctic region. But, there were 79 species only distributed in the Palaearctic region, accounting for 64.75% of the total known number. And other species were distributed by cross- regions. For example, there were 22 species distributed across the Palaearctic and Nearctic regions; 15 species across the Palaearctic and Oriental regions. Table 1 showed the species diversity composition and their zoogeographical distribution.

3.2. Ecological properties analysis to the Miridae of Hulun Buir city

Because vegetation with a clear geographical differentiation in the horizontal direction in Hulun Buir city [6], it is necessary to study horizontal distribution of Miridae. The latitudinal range (47°47'-52°04' N) was further divided into 6 intervals. Our results showed that abundant species were distributed in the two intervals of 48°01'-49°N and 49°01'-50°N. Deraeocorinae had the largest species number (8 species) at the interval of 48°01'-49°N; while Mirinae (41 species), Orthotylinae (13 species) and Phylinae (10 species) all reached the highest at the interval of 49°01'-50°N. Details of horizontal distribution of Miridae were shown in Fig. 1.




Fig. 1. Horizontal distribution of Miridae from Hulun Buir city

Our data also revealed the range of vertical distribution of Miridae (Fig. 2). We divided the whole altitudinal range (318-834m) into 5 intervals. For the family Miridae, the abundance reached a peak (86 species) in 601-800m which took 70.49 % of the total species number; there was no distributional record in 0-200 m. From the perspective of the subfamily, Mirinae, Orthotylinae and Phylinae all reached the highest species number (49 species, 16 species, 18 species) within the range of 601-800 m; while the Deraeocorinae subfamily reached the highest species number within the range of 201-400 m. This circumstance suggested that the species of Deraeocorinae was more suitable for living in low altitude area than other subfamilies.




Fig. 2. Vertical distribution of Miridae from Hulun Buir city

4. Conclusion and Discussion

Our study showed a total of 122 species of Miridae from Hulun Buir city, distributed in four animal geographical regions. The Palearctic composition was dominant. Studies of ecological properties had shown about half of the species were distributed in both latitudinal interval 48°01'-49°N and 49°01'-50°N. While in the range of 48°01'-50°N was just covered the Hulun Buir Grassland. This also confirmed that Miridae was very suitable for the survival in the grassland. In addition, the most appropriate altitudinal range was 601-800 m. Our results also suggested the adaptability of Mirinae to the environment was stronger; while Deraeocorinae was more suitable for low latitude and low altitude area than other subfamilies. For the Miridae possessed above-mentioned species diversity and ecological properties characteristics, the reason

probably was its special geographical location and overall environmental characteristics and other factors. Our study reflects the basic situation of Miridae from Hulun Buir city. But the city is very vast, the habitat, the climate, the vegetation type is also complex and diverse. At present, the research on Miridae is still very weak, on the insect's further biological geography, biology and ecology research needs to strengthen.

5. Acknowledgements

The research work was supported by the funds of the National Natural Science Foundation of China (31301922).

6. References

- [1] R. Liu, M. Feng, J. L. Sun, *et al.* The vegetation classification based on HJ-CCD data and decision tree. *Sci.Geog. Sin.*, 2012, **32**(12):1488-1495.
- [2] I. M. Kerzhner. Key to the Insects of the Far East of the USSR. In: Lehr P. A. (Ed.), *Family Miridae*. Vol. 2. Leningrad: Nauka. 1988, pp.778-857.
- [3] I. M. Kerzhner, M. Josifov. Catalogue of the Heteroptera of the Palaearctic Region. In: Aukema B., Christian R. (Eds.), *Cimicomorpha II*, Vol. 3. Amsterdam: The Netherlands Entomological Society. 1999, pp. 204-277.
- [4] N. J. Tatarnic, G. Cassis. The Halticini of the world (Insecta: Heteroptera: Miridae: Orthotylinae): generic reclassification, phylogeny, and host plant associations. *Journ. of the Linnean Soc. Zool.*, 2012, **164**(3): 558-658.
- [5] A. R. Wallace. *The geographical distribution of animals*. New York: Harper & Brothers, 1876.
- [6] H. S. Wu, Nonnaizab. *Grasshoppers of Hulunbeier Grassland*. Beijing: China Agricultural Press, 2009.